

# Update on DUNE physics

IN2P3 Scientific Council 2-3-2026


A.Meregaglia (LP2i Bordeaux)

# Introduction

**Neutrino: one of the least known elementary particles**

## What we know

- Neutrinos of **3 families** have been detected and **oscillations** between the families were observed.
- The three **mixing angles** and the two **mass splittings** have been measured with precision at the **few percent level**.
- **Neutrino masses** are the first and so far only manifestation of physics **beyond the Standard Model** in particle physics experiments.



NO Param	Ref. [181] w/o SK-ATM Best Fit Ordering	
	bfp $\pm 1\sigma$	$3\sigma$ range
$\frac{\sin^2 \theta_{12}}{10^{-1}}$	$3.03^{+0.12}_{-0.11}$	2.70 $\rightarrow$ 3.41
$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	31.31 $\rightarrow$ 35.74
$\frac{\sin^2 \theta_{23}}{10^{-1}}$	$5.72^{+0.18}_{-0.23}$	4.06 $\rightarrow$ 6.20
$\theta_{23}/^\circ$	$49.1^{+1.0}_{-1.3}$	39.6 $\rightarrow$ 51.9
$\frac{\sin^2 \theta_{13}}{10^{-2}}$	$2.203^{+0.056}_{-0.059}$	2.029 $\rightarrow$ 2.391
$\theta_{13}/^\circ$	$8.54^{+0.11}_{-0.12}$	8.19 $\rightarrow$ 8.89
$\delta_{CP}/^\circ$	$197^{+42}_{-25}$	108 $\rightarrow$ 404
$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	6.82 $\rightarrow$ 8.03
$\frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2}$	$2.437^{+0.028}_{-0.027}$	2.354 $\rightarrow$ 2.523
IO	$\Delta\chi^2 = 2.3$	
$\frac{\sin^2 \theta_{12}}{10^{-1}}$	$3.03^{+0.12}_{-0.11}$	2.70 $\rightarrow$ 3.41
$\theta_{12}/^\circ$	$33.41^{+0.75}_{-0.72}$	31.31 $\rightarrow$ 35.74
$\frac{\sin^2 \theta_{23}}{10^{-1}}$	$5.78^{+0.16}_{-0.21}$	4.12 $\rightarrow$ 6.23
$\theta_{23}/^\circ$	$49.5^{+0.9}_{-1.2}$	39.9 $\rightarrow$ 52.1
$\frac{\sin^2 \theta_{13}}{10^{-2}}$	$2.219^{+0.060}_{-0.057}$	2.047 $\rightarrow$ 2.396
$\theta_{13}/^\circ$	$8.57^{+0.12}_{-0.11}$	8.23 $\rightarrow$ 8.90
$\delta_{CP}/^\circ$	$286^{+27}_{-32}$	192 $\rightarrow$ 360
$\frac{\Delta m_{21}^2}{10^{-5} \text{ eV}^2}$	$7.41^{+0.21}_{-0.20}$	6.82 $\rightarrow$ 8.03
$\frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2}$	$-2.498^{+0.032}_{-0.025}$	-2.581 $\rightarrow$ -2.408


from PDG (2023)

# Introduction

Neutrino: one of the least known elementary particles

## What we do not know

- The **absolute** value of their **masses**.
- Whether neutrinos are **Dirac or Majorana fermions**.
- The **mass ordering** (normal or inverted).
- Whether the **charge-parity (CP) symmetry** is violated in the lepton sector.
- Whether the **three-neutrino framework** is complete and the PMNS matrix is unitary.




# Introduction

Neutrino: one of the least known elementary particles

## What we do not know

- The **absolute** value of their **masses**.
- Whether neutrinos are **Dirac or Majorana fermions**.
- The **mass ordering** (normal or inverted).
- Whether the **charge-parity (CP) symmetry** is violated in the lepton sector.
- Whether the **three-neutrino framework** is complete and the PMNS matrix is unitary.




Likely measured in the next 10 years with critical contributions from Long Baseline experiments

# Long Baseline Experiments


- The precise study of oscillations with Long Baselines (LBL) is a **privileged tool** to investigate both standard and non-standard neutrino properties.
- The **main goals** of current and future LBL experiments are the **search for CPV** in the neutrino sector, the **determination of the neutrino mass ordering**, and the **precise measurement** of the phase  $\delta_{CP}$  and of the so called “atmospheric” parameters ( $\theta_{23}$  and  $\Delta m^2_{32}$ ).
- T2K and NOVA are providing accurate measurements of neutrino oscillations and new tests of the standard PMNS oscillation paradigm, observed for the first time  $\nu_e$  appearance in a  $\nu_\mu$  beam, and yielded first hints of CP-symmetry violation.
- The **next generation experiments**, Hyper-Kamiokande (HK) in Japan and the Deep Underground Neutrino Experiment (**DUNE**) in the US, represent the **forefront of this endeavour**.

# DUNE


- DUNE is a **long baseline neutrino oscillation experiment** with a baseline of 1300 km.
- The neutrino beam will be produced at **Fermilab** and detected in the **SURF** (Sanford Underground Research Laboratory) underground facility (1500 m overburden).
- The experiment consists of a 1.2 MW proton beam upgradable to more than 2 MW, a near detector complex (at about 570 m from the target), and more than **40 kton far detector** (staged approach).





## FD complex



## ND complex





# DUNE FD technology



- The liquid Argon Time Projection Chamber (**LArTPC**) was proposed by C.Rubbia in 1977, and it is a homogeneous massive target and ionization detector, i.e. an **electronic bubble chamber**.
- Ideal detector for neutrino oscillations, supernovae neutrinos and proton decay search:
  - Large mass, homogeneous detector, low thresholds, exclusive final states
  - Tracking + calorimetry (0.035  $X_0$  sampling with 5mm pitch)
  - Electron identification,  $\pi^0$  rejection, particles identification with dE/dx

# DUNE FD technology




**More on DUNE FD in Cédric's talk**

- The Liquid Argon Projection Chamber (**LArTPC**) was proposed by C.Rubbia in 1977, as a homogeneous massive target and ionization detector, i.e. an **electronic chamber**.
- Ideal detector for neutrino oscillations, supernovae neutrinos and proton decay search:
  - Large mass, homogeneous detector, low thresholds, exclusive final states
  - Tracking + calorimetry ( $0.035 X_0$  sampling with 5mm pitch)
  - Electron identification,  $\pi^0$  rejection, particles identification with  $dE/dx$

# DUNE ND

- DUNE will have a **Near Detector complex critical starting from early physics goals** to reduce flux and cross sections uncertainties and have the most accurate prediction of neutrino spectrum at the Far Detector.
- **System for on-Axis Neutrino Detection (SAND):** On-axis magnetized low-density tracker and calorimeter to control and monitor the neutrino beam on axis.
- **Near Detector Liquid Argon (NDLAr):** Liquid Argon Detector (50t fiducial) with the same target and technology as FD.
- **The Muon Spectrometer (TMS):** detector to measure muons escaping from NDLAr.
- **NDLAr+TMS** are a **movable** detector and exploit the **PRISM technique**: they measure neutrino flux at different angles (i.e. different energies) for a **better extrapolation at the FD reducing the dependency on the accuracy of interaction models**.



# DUNE staged approach

- DUNE follows a **staged approach** which implies the construction of different ND and FD modules and beam upgrades which have been **updated following the P5 report** in 2023 and is made of two phases.

## Phase 1

### **2026 start detector installation - 2029 physics - 2032 beam + ND**

- Full near + far site facility and infrastructure.
- Two LArTPC modules (total active mass of about 25 kton).
- Upgradeable 1.2 MW neutrino beamline.
- Movable LArTPC near detector and on axis detector (NDLAr + TMS + SAND).


## Phase 2

### **More detectors and beam upgrade**

- Two additional FD modules (more than 40 kton fiducial in total).
- Beamline upgrade to multi MW.
- More capable near detector (including ND-GAr).

# DUNE oscillation strategy

- DUNE will measure neutrino spectra in a **wide band beam exploiting the spectral information** and 2<sup>nd</sup> oscillation maximum.
- DUNE will study  $\nu_\mu \rightarrow \nu_e$  oscillations both with neutrinos and antineutrinos.




- Wide band beam and long baselines **break the degeneracy between  $\delta_{CP}$  and matter effects**.
- **DUNE is a unique experiment** enabling searches for CP violation, mass ordering, and three-flavour neutrino phenomenology, including  $\nu_\tau$  appearance, within a single experimental setup, while providing detailed reconstruction of neutrino interaction final states.

# CP violation - general considerations

- Typically to measure CP violation the idea is to measure and compare  $\nu_\mu \rightarrow \nu_e$  and  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  transitions.

## Few important considerations

- **Matter effects and CP are degenerate** since they both impact neutrino and antineutrino in opposite ways, however matter effects depends on the ordering whereas CP effects do not.
- **Narrow band** experiments basically just “count” the appeared neutrinos.
- There are therefore solutions with CP conserving phases in one ordering which overlap to CP violating phases in the opposite ordering.




Maximal CP violation IO

CP conserving NO

# CP violation - HK




- The next important actor in the search of CP violation will be Hyper-Kamiokande (HK) starting in 2028.
- The experiment uses the same technology as T2K but with an upgraded beamline (factor of about 2 with respect to current beam power of 750 kW) and an increase in mass of a factor of 8.
- The bi-plot is the same with a smaller error.
- HK is excellent at distinguishing between conserving and violating CP phases **provided the mass ordering is known.**



# CP violation - DUNE strengths

- The specific **advantages of DUNE** in the search of CP violations are:
  - ➔ **Higher energy** (to match the longer baseline) with **larger matter effects**.
  - ➔ The use of a **wide band beam** (WBB) which allows for **spectral information** exploitation.

**No spectral information**




**The NH and IH ellipses are well separated and no degeneracy remains**


# CP violation - DUNE strengths

## Spectral information



### High L/E



### Medium L/E




### Low L/E




- DUNE can measure oscillation parameters in **several regions of L/E**.
- If 3-flavor PMNS picture is correct, data should prefer the same point in every bin.
- It is likely that **new physics** would distort the result as a function of L/E (DUNE unique sensitivity).


# CP violation - Sensitivity



7 years staged  
10 years staged  
15 years staged




Best scenario




**DUNE could have  $>3 \sigma$  CPV sensitivity in 3.5 years**

# CP violation - Sensitivity




7 years staged  
10 years staged  
15 years staged

Best scenario






*Expected question... how does this compare to HK?*

*DUNE could have  $>3 \sigma$  CPV sensitivity in 3.5 years*




# CP violation - Sensitivity

**Comparable results  
between DUNE and HK**



**DUNE could have the best  
ultimate  $\delta_{CP}$  resolution,  
especially if CP is violated.**




# Mass ordering - general considerations

- There are several ways to tackle the mass ordering:


- **Atmospheric neutrinos:** exploit measurements over a wide range of energies and baselines (zenith angles), comparing the observed patterns with expectations for normal and inverted ordering.

e.g. SK, ORCA



- **Reactor neutrinos:** measure with very high precision the frequency of the wiggles related to  $\Delta m^2_{31}$  in  $\bar{\nu}_e$  oscillations.

e.g. JUNO





- **Long-baseline accelerator experiments:** exploit the enhancement or suppression of  $\nu_e$  appearance induced by matter effects.

e.g. DUNE



# Mass ordering - DUNE

- DUNE is the **only experiment** that measures the mass ordering by **separately** measuring  $\nu_\mu \rightarrow \nu_e$  and  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  oscillations.
- The exploitation of the spectral information is crucial.

Neutrinos



Antineutrinos



Statistics corresponding to an excess of about  $5\sigma$



40-50% differences in 0.5-2 GeV region, with opposite sign in FHC and RHC

Worst scenario with CPV and matter effect in opposite directions

# Mass ordering - DUNE


- Although DUNE's mass ordering sensitivity depends on true oscillation parameters, **DUNE will measure mass ordering no matter what.**
- In the **favourable** scenario, DUNE surpasses  **$5\sigma$**  in less than 1 year, but even in the most **pessimistic** scenario, DUNE reaches **more than  $5\sigma$**  in about 5 years.

**Although starting later DUNE is the only experiment that will reach  $5\sigma$  without combination with other experiments and regardless of the true ordering.**



# Solar neutrino

- The possibility to study solar neutrinos in DUNE has been boosted by the work on detecting low energy events as demonstrated with radiological events in ProtoDUNE.
- Separate precision measurements of neutrino-mixing parameters and the  $^8\text{B}$  flux could be made using two channels and the day-night effect.




**Best precision on solar given by DUNE alone with two caveat: a threshold at 5 MeV and a shielding to reduce low energy background (work in progress)**

**CC channel:**  $\nu_e + {}^{40}\text{Ar} \rightarrow e^- + {}^{40}\text{K}^*$

**ES channel:**  $\nu_x + e^- \rightarrow \nu_x + e^-$  (pointing)



	10 MeV Threshold	5 MeV Threshold
<b>CC</b>	1300	117000
<b>ES</b>	1400	33000



# 3 flavour paradigm

## oscillation parameters

- DUNE will have a unique opportunity to test the **PMNS unitarity** and confirm the three flavour paradigm.
- Measurements  $\theta_{13}$  obtained from long-baseline ( $\nu_e$  appearance) and reactor experiments ( $\bar{\nu}_e$  survival) are only equivalent if unitarity is assumed.




**DUNE approaching world leading measurements on  $\theta_{13}$  and  $\Delta m^2_{atm}$**

# 3 flavour paradigm


$$\nu_\mu \rightarrow \nu_\tau$$

- Another opportunity of DUNE to test unitarity is directly **measure 3-flavor oscillations**.
- Thanks to the high-energy proton beam with on-axis detector, DUNE has substantial flux above the 3.5 GeV  $\nu_\tau$  CC threshold and could therefore **observe  $\nu_\tau$  appearance**.
- Imaging capabilities of DUNE's LArTPCs makes it possible to **reconstruct complex  $\tau$  final states kinematically**.
- ML-driven reconstruction based on hierarchical graph neural network being pursued.





# Atmospheric neutrinos

- Atmospheric neutrinos will be **first DUNE data** (between 2029 and 2032) and will be important for commissioning the detector and for physics studies.





- DUNE analysis in progress, with joint fit framework in place.
- Sub-GeV neutrinos could **exclude several values of CP** phases beyond  $3\sigma$ .
- GeV neutrino could give a first measurement of **mass ordering complementary to beam** measurements.




# BSM

- DUNE is sensitive to some specific oscillation modes.


- Sterile neutrinos** can be tackled thanks to the large range of  $L/E$  (wide-band beam and ND/FD and atmospheric neutrinos).



- Precise measurements of neutrinos and antineutrinos could give the **best limits on CPT violation**.



- Large matter effects and three-flavour sensitivity allow DUNE to search for **non-standard interactions**.



# Supernova





- Time and energy profile of SNB neutrino flux is rich in supernova astrophysics.

- Flux includes all neutrino flavours but **only DUNE measures  $\nu_e$**  thanks to the Ar target.

- DUNE is uniquely sensitive to mass ordering via SNB detection.

- About  $5^\circ$  pointing resolution thanks to ES channel depending on SNB location (40 kton 10 kpc)



- DUNE supernova program starts at the moment FD1 is turned on. Reach extends beyond the Milky Way with additional modules.



	$\nu_e$	$\bar{\nu}_e$	$\nu_x$
DUNE	89%	4%	7%
SK <sup>1</sup>	10%	87%	3%
JUNO <sup>2</sup>	1%	72%	27%

<sup>1</sup>Super-Kamiokande, *Astropart. Phys.* **81** 39-48 (2016)

<sup>2</sup>Lu, Li, and Zhou, *Phys Rev. D* **94** 023006 (2016)



# Proton decay

- DUNE will be an **excellent detector to perform nucleon decay searches** considering the underground location, the very large fiducial mass, and the millimetric size imaging capabilities.

- The **golden channel** would be





The main background for this channel in DUNE are atmospheric neutrinos.

The 90% CL lower limit (400 kton-year exposure) is

$$\tau/\text{Br}(p \rightarrow K^+ + \bar{\nu}) > 1.3 \times 10^{34} \text{ years}$$

**DUNE sensitivity comparable to larger water Cherenkov detectors**



# IN2P3 contributions

- There is a **large implication of IN2P3 groups in various critical topics.**

## Physics analysis

- APC → Atmospheric, sterile and supernova neutrinos, and systematics studies.
- LAPP → Solar neutrinos.
- IJCLab → Low energy events.
- IP2I → LBL neutrinos and energy reconstructions.
- LP2iB → LBL neutrinos and energy reconstructions.
- LPSC → LBL sensitivity, low energy showers (GeV), PID and ML.

## CERN prototypes analysis


- APC → Neutrino events, pion energy reconstruction, and PDS related analyses.
- LAPP → Data reconstruction and analysis leader.
- IJCLab → Data reconstruction (incl. PDS), processing & simulation, detector calibration.
- IP2I → Online analysis and hadronic interactions.
- LP2iB → Validation of data/MC response.
- LPSC → CRP related studies.

## Computing/software


- IP2I → Management of data at CC Lyon and responsible in the computing consortium for data and MC production.

# Energy resolution/LBL oscillations


- Work ongoing to understand the best way to reconstruct EM showers (*IP2I and LP2I*).
- Work started in parallel to select showers in PDVD data and improve PID in general with the DUNE AI/ML workflow SPINE (*IJCLab and LPSC*).
- Deployment on LPSC HPC platform of a framework to estimate DUNE LBL sensitivity implementing oscillation analysis with GUNDAM with updated systematics and VD performances (*LPSC*).
- Study of the impact of CRP gaps, implementation of correct geometry and tuning of charge recovering based on PDVD data (*LP2I*).



From IP2I



From LPSC




From LP2I

# Atmospheric neutrinos

- The study of atmospheric neutrinos is a primary topic for DUNE and France is well visible and recognized on this (*effort driven by APC*).
- The **first paper on atmospheric neutrino reconstruction** has been released [[arXiv:2601.05697](https://arxiv.org/abs/2601.05697)] (co-lead by *APC*).
- In addition studies on **sterile neutrino searches** were performed (*APC*).
- The study of atmospheric neutrinos includes the reconstruction of vertex (*APC*) with impacts also on beam events .


- The identification of neutrinos and antineutrinos via the Michel electron tagging is under study to improve the sensitivity to mass ordering (*APC*).



From [arXiv:2601.05697](https://arxiv.org/abs/2601.05697)


# Solar neutrinos

- The study of solar neutrino has also started in France, initiated at *LAPP*.
- The evaluation of background in particular neutrons and cosmogenic backgrounds was carried out and the sensitivity estimated in the current situation and with shielding (*LAPP*).




From *LAPP*

- Calibration at low energy is critical for solar neutrinos but also for SuperNova neutrinos. This work is ongoing in connection with the calibration team, using radiologicals (*IJCLab and LAPP*), and Michel electrons (*IJCLab*).



From *IJCLab*





From *IJCLab*

# Systematics for atmospheric and LBL

- The assessment of systematic is a topic transverse to different WGs (LBL, ANOA, FD Sim/Reco, Calibration) on which France has always been involved (*effort driven by APC*).
- A new framework for FD systematics has been developed (presented@LBNC in October 2025) and is being improved (*APC*).
- The beam data on ProtoDUNE will be analyzed to evaluate systematics on energy reconstruction (*APC*).


## Framework for FD systematics studies, ANOA+LBL




From APC

# ProtoDUNE and Coldbox analysis

- France is heavily involved in ProtoDUNE, co-leader of the Data Reconstruction and Analysis working group, and development of dedicated tools (LARDON) systematically used during prototyping (*LAPP*).
- A lot of work is done to assure the correct data readability and utilization (*IP2I*).
- France is also involved in the Data Production (*IJCLab and IP2I*).
- Analysis performed in the VD coldbox (PDS, charge) will be extended to PDVD (*APC and LAPP*).
- Work on the transparency of CRPs as a function of the voltage and the reconstructed charge as a function of the track angle (*LAPP and LPSC*), will be validated on PDVD.
- First studies on PDVD and PDHD data were presented (*IJCLab and LAPP*).
- First neutrino-Argon interactions seen in DUNE (*APC*).



From APC



From APC (LARDON display)


# Conclusions

- The DUNE physics case, already assessed in previous CS in 2018 and 2021. Its importance was restated and **supported by the new P5 and European Strategy outcomes**.
- DUNE is designed to **test the three-flavor paradigm** and search for new physics in neutrino oscillations.
- If the three-flavor picture is correct, DUNE will make **very precise measurements of its parameters**, leveraging spectral information to break degeneracies.
- DUNE has qualitatively **superior ability to determine the neutrino mass ordering** definitively.
- DUNE has unique capabilities to run a **rich physics program without beam**.
- There is a **large implications of IN2P3 groups in the physics analysis, analysis of CERN prototypes** (only real data before first FD operation) and **computing/software efforts**.
- Strong IN2P3 efforts for a timely operation of the first FD module (see Cédric's talk).

# BackUp

# ACE MIRT

- An increase in **beam power to > 2 MW is a primary requirement for DUNE** to achieve its P5 physics goals.
- Fermilab has in place an **Accelerator Complex Evolution (ACE)** plan to realize the beam power required. It will be delivered in two phases:
  1. The first phase known as **ACE-Main Injector Ramp and Targetry (ACE-MIRT)**, will raise the beam power to just beyond 2 MW by reducing the cycling time in the Main Injector (MI) from 1.2 sec to 0.7 sec (available since 2031 !)
  2. Phase two is the **replacement of the Booster and an extension of the Linac to 2 GeV (ACE-BR)**, which will eventually allow the beam power to rise to 2.4 MW.
- While the ACE-MIRT upgrade is primarily motivated by DUNE science, ACE-BR targets a broader physics program at FNAL, serving as a robust and reliable platform for the future.



# Octant degeneracy

- The  $\theta_{23}$  angle is currently known with  $\sin^2\theta_{23}$  roughly between 0.41 and 0.62 at  $3\sigma$ .
- DUNE can measure  $\sin^2\theta_{23}$  with  $3\sigma$  precision in the range  $\sim 0.48\text{--}0.55$ , allowing a determination of the  $\theta_{23}$  **octant**.
- $\nu_{\mu} \rightarrow \nu_{\mu}$  disappearance depends on  $\sin^2 2\theta_{23}$ , while  $\nu_{\mu} \rightarrow \nu_e$  appearance depends on  $\sin^2\theta_{23}$ ; combining both channels helps resolve the octant degeneracy.

